

Name:

SUMMER 2017 — MATH 110
FINAL EXAM

Section	Max	Score
Part I — True/False	15	
Part II — Examples	10	
Part III — Calculations	20	
Part IV — Proofs	10	
Total	55	

Part I — True/False. You will get 1 point for each correct answer, 0 points for each blank answer, and −1 point for each incorrect answer. The minimum possible score for this section is 0.

- (1) Let $V = \{(a, b) : a, b \in \mathbf{R}\}$. Define addition on V coordinate-wise, and define a scalar multiplication operation $@$ by the formula

$$\lambda @ (a, b) = (0, \lambda b)$$

for all $(a, b) \in V$ and $\lambda \in \mathbf{R}$. Then V , equipped with these operations, is a vector space over \mathbf{R} .

- (2) The set $\{(a, b, c) \in \mathbf{R}^3 : a^3 = b^3\}$ is a subspace of \mathbf{R}^3 . T F
- (3) Suppose p_0, p_1, p_2 is a list of four polynomials in $\mathcal{P}_2(\mathbf{R})$ with the property that $p_i(1) = 0$ for each i . Then the list p_0, p_1, p_2 is linearly dependent. T F
- (4) Suppose U is a subspace of a vector space V and define a map $T : V \rightarrow V$ by T F

$$T(v) = \begin{cases} v & \text{if } v \in U \\ 0 & \text{if } v \notin U. \end{cases}$$

Then T is linear.

- (5) Every basis for $\mathcal{P}_3(\mathbf{F})$ contains a degree 2 polynomial. T F
- (6) Suppose v_1, v_2, v_3 is a basis for V and U is a subspace of V such that $v_1 \in U$ but $v_2, v_3 \notin U$. Then $U = \text{span}(v_1)$. T F
- (7) If V and W are vector spaces, $T \in \mathcal{L}(V, W)$, and v_1, \dots, v_n is a list in V such that Tv_1, \dots, Tv_n is linearly independent, then v_1, \dots, v_n is linearly independent. T F
- (8) Suppose V is a finite dimensional inner product space. Every self-adjoint $T \in \mathcal{L}(V)$ has a cube root. T F
- (9) There exists $T \in \mathcal{L}(\mathbf{R}^4, \mathbf{R}^2)$ such that $\text{null } T = \{(0, 0, 0, x) : x \in \mathbf{R}\}$. T F
- (10) Suppose W_1, W_2 and U are all subspaces of a vector space V such that $U \oplus W_1 = V$ and $U \oplus W_2 = V$. Then $W_1 = W_2$. T F
- (11) Suppose V is a finite dimensional inner product space, $U = \text{span}(u_1, u_2, u_3)$ is a subspace, and $v \in V$ is a vector such that T F

$$|\langle v, u_1 \rangle|^2 + |\langle v, u_2 \rangle|^2 + |\langle v, u_3 \rangle|^2 = 0.$$

Then $v \in E(1, P_{U^\perp})$.

- (12) Suppose $T \in \mathcal{L}(\mathbf{R}^4)$ is not injective and $\dim E(9, T) = 3$. Then T is diagonalizable. T F
- (13) Suppose $T \in \mathcal{L}(\mathbf{C}^4)$ satisfies $(T - 3I)(T - 2I)(T - I) = 0$. Then the minimal polynomial p_{\min} of T is $p_{\min}(z) = (z - 3)(z - 2)(z - 1)$. T F
- (14) Suppose V is a vector space, $T \in \mathcal{L}(V)$ is an operator, and $v \in V$ is a vector such that $(T - 3I)(T + 3I)v = 0$. Then v is an eigenvector of T , with eigenvalue either 3 or −3. T F
- (15) Suppose V is the inner product space of continuous functions $[-1, 1] \rightarrow \mathbf{R}$, with inner product given by T F

$$\langle f, g \rangle = \int_{-1}^1 f(x)g(x) dx.$$

Let $h \in V$ be given by $h(x) = |x|$. Then $P_{\text{span}(1, x, x^2)}(h) = P_{\text{span}(1, x^2)}(h)$.

Part II — Examples. For each of the following, give an example of an operator $T \in \mathcal{L}(\mathbf{C}^4)$ that has the required properties by writing down a matrix *in Jordan form* that represents the operator, if it is possible. If it is impossible, write “impossible.” A correct answer is worth 2 points, a blank answer is 0.5 points, and an incorrect answer is 0 points. No justification is required.

(1) T has precisely 3 distinct eigenvalues and its minimal polynomial is of degree 3.

(2) T is nilpotent and there does *not* exist a basis of \mathbf{C}^4 of the form T^3v, T^2v, Tv, v for some $v \in \mathbf{C}^4$.

(3) $\dim \text{null}(T - 2I) = 2$, $\dim \text{null}(T - 2I)^2 = 3$ and $\dim \text{null}(T - 2I)^3 = 4$.

(4) There is no 3 dimensional subspace of \mathbf{C}^4 invariant under T .

(5) The quotient operator T/U is not injective, where $U = \text{null } T$.

Part III — Calculations. In each of the following, you are asked to calculate the dimension of a vector space. Write the dimension inside the box on the right. If the dimension is infinite, write “ ∞ .” A correct answer is worth 2 points, a blank answer is 0.5 points, and an incorrect answer is 0 points. No justification is required.

(1) Let $U = \{(x, y, z) \in \mathbf{R}^3 : x + 2y + 3z = 0\}$. Calculate $\dim(\mathbf{R}^3/U)$.	
(2) Calculate $\dim \mathcal{L}(\mathcal{P}_2(\mathbf{R})) \times \mathcal{P}_3(\mathbf{R})'$.	
(3) Let $T \in \mathcal{L}(\mathcal{P}_4(\mathbf{R}), \mathcal{P}_3(\mathbf{R}))$ be defined by $T(f)(z) = zf''(z)$. Calculate $\dim \text{range } T$.	
(4) Suppose $T \in \mathcal{L}(\mathbf{C}^4)$ has 4 distinct eigenvalues and $p_{\text{char}}(0) = 0$. Calculate $\dim \text{range } T$.	
(5) Suppose U is the space of self-adjoint operators inside $\mathcal{L}(\mathbf{R}^3)$. Calculate $\dim U$.	
(6) Suppose $T \in \mathcal{L}(\mathbf{R}^5, \mathbf{R}^3)$ and $\dim \text{null } T = 2$. Calculate $\dim \text{null } T'$.	
(7) Suppose $T \in \mathcal{L}(\mathbf{F}^\infty)$ is the forward shift operator $T(x_0, x_1, x_2, \dots) = (0, x_0, x_1, \dots)$ and define $U = \text{range } T$. Calculate $\dim \mathbf{F}^\infty/U$.	
(8) Suppose V is the vector space of infinitely differentiable functions $\mathbf{R} \rightarrow \mathbf{R}$ and $T \in \mathcal{L}(V)$ is the operator $T(f) = f' - f$. Calculate $\dim \text{null } T$.	
(9) Suppose $S, T \in \mathcal{L}(\mathbf{R}^3, \mathbf{R})$ are nonzero linear maps with distinct null spaces. Calculate $\dim(\text{null } S) \cap (\text{null } T)$.	
(10) Regard \mathbf{C}^2 as a <i>real</i> vector space and \mathbf{R}^2 as a subspace of \mathbf{C}^2 . Calculate $\dim(\mathbf{C}^2/\mathbf{R}^2)$.	

Part IV — Proofs. You must write rigorous arguments using complete sentences for full credit. Each problem is worth a maximum of 5 points.

(1) For an arbitrary vector space V , prove that $\mathcal{L}(\mathbf{F}^2, V)$ is isomorphic to $V \times V$.

(2) Suppose $T \in \mathcal{L}(\mathcal{P}(\mathbf{F}))$ is injective and $\deg Tp \leq \deg p$ for all $p \in \mathcal{P}(\mathbf{F})$. Prove that T is invertible.

